

Minimal hypersurfaces in $\mathbb{S}^5$ with vanishing Gauss-Kronecker curvature

Marcos M. Diniz, José Antonio M. Vilhena and Juan F. Z. Zapata

Abstract. In this paper we present a local description for complete minimal hypersurfaces in $\mathbb{S}^5$ with zero Gauss-Kronecker curvature, zero 3-mean curvature and nowhere zero second fundamental form.

Mathematics Subject Classification (2010). 53C42.

Keywords. Minimal hypersurfaces, Gauss-Kronecker curvature, complete hypersurfaces.

Introduction

Let $\mathbf{g} : M^2 \hookrightarrow \mathbb{S}^4$ be a minimal immersion, where M^2 is a 2-dimensional manifold. Dajczer and Gromoll in [2] proved that if $\mathbf{g}$ has nowhere vanishing normal curvature, then the polar map of $\mathbf{g}$ is everywhere regular and provides a minimal hypersurface in $\mathbb{S}^4$ with Gauss-Kronecker curvature identically zero. In [3] de Almeida and Brito classified the compact minimal hypersurfaces M^3 in $\mathbb{S}^4$ with zero Gauss-Kronecker curvature and non-vanishing second fundamental form on M^3 . Later, Ramanathan [8] classified the compact minimal hypersurfaces M^3 in $\mathbb{S}^4$ with zero Gauss-Kronecker curvature without the condition on the second fundamental form. In [5] Hasanis, Halilaj and Vlachos classified the complete minimal imersions $\mathbf{g} : M^2 \hookrightarrow \mathbb{S}^4$ with Gauss-Kronecker curvature identically zero under some assumptions on the second fundamental form. In [1] Asperti, Chaves and Sousa Jr. showed that the infimum of the absolute value of the Gauss-Kronecker curvature of a complete minimal hypersurfaces in 4-dimensional space form vanishes under assumptions on the Ricci curvature and classified the complete minimal

CAPES - PROCAD-NF/2010.

CAPES - PROCAD-NF/2010.

CNPq - BOLSA

August 02, 2012.

hypersurfaces in 4-dimensional space form with Gauss-Kronecker curvature constant.

In this paper, we consider $\mathbf{g}: M^2 \hookrightarrow \mathbb{S}^5$ be a minimal immersion, where M^2 is a complete 2-dimensional manifold. Denote by K and K_N (see (1.12)) the Gauss and scalar normal curvature, respectively. In Proposition 2.1 we show that $\mathbf{g}$ is superminimal if and only if

$$(K-1)^2 - \frac{1}{4} K_N = 0.$$

In Theorem 2.2 we give an example of a minimal hypersurface in $\mathbb{S}^5$ with vanishing Gauss-Kronecker and 3-mean curvatures. This example is a 2-spherical local bundle over a minimal surface in $\mathbb{S}^5$. Finally in Theorem 3.3 we study locally the complete minimal hypersurfaces in $\mathbb{S}^5$ with zero Gauss-Kronecker curvature, zero 3-mean curvature and nowhere zero second fundamental form.

1. Preliminaries

Isometric immersions in Euclidean sphere

Let $\mathbf{g}: M^n \hookrightarrow \mathbb{S}^{n+p}$ an isometric immersion from an n -manifold M^n into the $(n+p)$ -sphere. Let $\mathcal{B} = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n+p}\}$ be an orthonormal frame on $\mathbb{S}^{n+p}$ adapted to the immersion, in the sense that $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ spans $\mathbf{g}_*(TM)$, while $\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+p}\}$ spans $(\mathbf{g}_*(TM))^\perp$. Let $\mathcal{B}^* = \{\omega^1, \omega^2, \dots, \omega^{n+p}\}$ its dual co-frame. The structure equations of $g(M^n)$, in terms of this co-frame are given by

$$\left\{ \begin{array}{l} d\mathbf{g} = \omega^i \otimes \mathbf{e}_i; \\ d\mathbf{e}_i = -\omega^i \otimes \mathbf{g} + \omega_j^i \otimes \mathbf{e}_j + \omega_i^\alpha \otimes \mathbf{e}_\alpha; \\ d\mathbf{e}_\alpha = \omega_\alpha^j \otimes \mathbf{e}_j + \omega_\alpha^\beta \otimes \mathbf{e}_\beta; \\ d\omega^i = -\omega_j^i \wedge \omega^j, \quad \omega_j^i + \omega_i^j = 0; \\ d\omega^\alpha = -\omega_i^\alpha \wedge \omega^i, \end{array} \right. \quad (1.1)$$

where $1 \leq i, j \leq n$ and $n+1 \leq \alpha, \beta \leq n+p$. Throughout all this work, we shall use small Latin letters, $i, j, \dots$ (resp. Greek letters, $\alpha, \beta, \dots$) for indices which take values over the range $\{1, \dots, n\}$ (resp. $\{n+1, \dots, n+p\}$) and we use the Einstein convention for summation over crossed indices. The Gauss, Codazzi and Ricci equations are given, respectively, by

$$\left\{ \begin{array}{l} d\omega_j^i = -\omega_k^i \wedge \omega_j^k - \omega_\alpha^i \wedge \omega_j^\alpha + 1 \omega^i \wedge \omega^j; \\ d\omega_\alpha^i = -\omega_k^i \wedge \omega_\alpha^k - \omega_\beta^i \wedge \omega_\alpha^\beta; \\ d\omega_\beta^\alpha = -\omega_i^\alpha \wedge \omega_\beta^i - \omega_\gamma^\alpha \wedge \omega_\beta^\gamma. \end{array} \right. \quad (1.2)$$

So, the curvature form Ω_j^i and normal curvature form Ω_β^α are

$$\begin{cases} \Omega_j^i = -\omega_\alpha^i \wedge \omega_j^\alpha + 1 \omega^i \wedge \omega^j, & \Omega_j^i = \frac{1}{2} R_{jkl}^i \omega^k \wedge \omega^l; \\ \Omega_\beta^\alpha = -\omega_i^\alpha \wedge \omega_\beta^i, & \Omega_\beta^\alpha = \frac{1}{2} R_{\beta kl}^\alpha \omega^k \wedge \omega^l. \end{cases} \quad (1.3)$$

We restrict these forms to M^n . Then

$$\omega^\alpha = 0.$$

Since

$$0 = d\omega^\alpha = -\omega_i^\alpha \wedge \omega^i,$$

by Catan's lemma we may write

$$\omega_i^\alpha = h_{ij}^\alpha \omega^j, \quad h_{ij}^\alpha = h_{ji}^\alpha. \quad (1.4)$$

We call

$$\mathbf{B} = h_{ij}^\alpha \omega^i \otimes \omega^j \otimes \mathbf{e}_\alpha \quad (1.5)$$

the *second fundamental form* of the immersed manifold M^n . The *mean curvature vector* of M^n is given by

$$\mathbf{H} = \frac{1}{n} \sum_i h_{ii}^\alpha \mathbf{e}_\alpha. \quad (1.6)$$

An immersion is said to be *minimal* if its mean curvature vanishes identically, i.e., if $\sum_i h_{ii}^\alpha = 0$ for all α .

Hypersurfaces

In the special case of hypersurfaces in $\mathbb{S}^5$, ($n = 4, p = 1$), taking a local orthonormal frame such that

$$h_{ij}^5 = \lambda_i \delta_j^i,$$

then the second fundamental form writes

$$\mathbf{B} = \lambda_i \omega^i \otimes \omega^i \otimes \mathbf{e}_5,$$

the equation (1.4) is written

$$\omega_i^5 = \lambda_i \omega^i. \quad (1.7)$$

the equations (1.1) writes

$$\begin{cases} d\mathbf{g} = \omega^i \otimes \mathbf{e}_i; \\ d\mathbf{e}_i = -\omega^i \otimes \mathbf{g} + \omega_j^j \otimes \mathbf{e}_j + \lambda_i \omega^i \otimes \mathbf{e}_5; \\ d\mathbf{e}_5 = -\lambda_i \omega^i \otimes \mathbf{e}_i; \\ d\omega^i = -\omega_j^j \wedge \omega^j, \quad \omega_j^j + \omega_i^i = 0; \\ d\omega^5 = 0, \end{cases} \quad (1.8)$$

and the Gauss and Codazzi equations (1.2) become

$$\begin{cases} d\omega_j^i = -\omega_k^i \wedge \omega_j^k + (1 + \lambda_i \lambda_j) \omega^i \wedge \omega^j; \\ d\omega_i^5 = -\lambda_k \omega_k^i \wedge \omega^k. \end{cases} \quad (1.9)$$

The r -mean curvatures H_r of an immersion $\mathbf{g}: M^4 \hookrightarrow \mathbb{S}^5$, with principal curvatures $\lambda_1, \lambda_2, \lambda_3$ and λ_4 , are given by

$$H_r = \frac{1}{\binom{n}{k}} \sum_{i_1 < i_2 \dots < i_r} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_r},$$

viz.,

$$\begin{cases} H_1 = \frac{1}{4}(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4); \\ H_2 = \frac{1}{6}(\lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_1 \lambda_4 + \lambda_2 \lambda_3 + \lambda_2 \lambda_4 + \lambda_3 \lambda_4); \\ H_3 = \frac{1}{4}(\lambda_1 \lambda_2 \lambda_3 + \lambda_1 \lambda_2 \lambda_4 + \lambda_1 \lambda_3 \lambda_4 + \lambda_2 \lambda_3 \lambda_4); \\ H_4 = \lambda_1 \lambda_2 \lambda_3 \lambda_4. \end{cases} \quad (1.10)$$

Note that H_1 and H_4 are, respectively, the *mean curvature* and the *Gauss-Kronecker curvature* of the hypersurface.

Immersed surfaces

In the case of immersed surfaces in $\mathbb{S}^5$ ($n = 2, p = 3$), equations (1.3) and (1.4) yield that the Gaussian curvature of M^2 is given by

$$K = \Omega_2^1(\mathbf{e}_1, \mathbf{e}_2) = d\omega_2^1(\mathbf{e}_1, \mathbf{e}_2) = 1 + \sum_{\alpha=3}^5 \begin{vmatrix} h_{11}^\alpha & h_{12}^\alpha \\ h_{21}^\alpha & h_{22}^\alpha \end{vmatrix}. \quad (1.11)$$

The *scalar normal curvature* of the immersion $\mathbf{g}$, K_N , is the length of the normal curvature form,

$$K_N = \sum_{i,j,\alpha,\beta} (R_{\beta ij}^\alpha)^2, \quad R_{\beta ij}^\alpha = \sum_k \begin{vmatrix} h_{ki}^\alpha & h_{kj}^\alpha \\ h_{ki}^\beta & h_{kj}^\beta \end{vmatrix}. \quad (1.12)$$

Let S be the square of the length of the second fundamental form,

$$S = \sum (h_{ij}^\alpha)^2.$$

Note that if $\mathbf{g}$ is minimal, then

$$K = 1 - \sum_{\alpha} \left[(h_{11}^\alpha)^2 + (h_{12}^\alpha)^2 \right], \quad K_N = 8 \sum \left(h_{11}^\alpha h_{12}^\beta - h_{12}^\alpha h_{11}^\beta \right)^2$$

i.e.,

$$K = 1 - \frac{1}{2}S, \quad K_N = 4 \left[(R_{412}^3)^2 + (R_{512}^3)^2 + (R_{512}^4)^2 \right]. \quad (1.13)$$

The *curvature ellipse* $\mathcal{E}_p$ of $\mathbf{g}$ at p is the image of the unitary circle by the second fundamental form $\mathbf{B}$ of $\mathbf{g}$ at p :

$$\mathcal{E}_p = \{\mathbf{B}_p(X, X) \in (T_p M)^\perp : X \in T_p M, \|X\| = 1\}.$$

For $X = \cos \theta \mathbf{e}_1 + \sin \theta \mathbf{e}_2$, it is easy to see that

$$\mathbf{B}_p(X, X) = \mathbf{H}_p + \begin{bmatrix} \mathbf{e}_3 & \mathbf{e}_4 & \mathbf{e}_5 \end{bmatrix} \cdot \begin{bmatrix} \frac{h_{11}^3 - h_{22}^3}{2} & h_{12}^3 \\ \frac{h_{11}^4 - h_{22}^4}{2} & h_{12}^4 \\ \frac{h_{11}^5 - h_{22}^5}{2} & h_{12}^5 \end{bmatrix} \cdot \begin{bmatrix} \cos 2\theta \\ \sin 2\theta \end{bmatrix}. \quad (1.14)$$

If $\mathbf{g}$ is minimal, then

$$\mathbf{B}_p(X, X) = \begin{bmatrix} \mathbf{e}_3 & \mathbf{e}_4 & \mathbf{e}_5 \end{bmatrix} \cdot \begin{bmatrix} h_{11}^3 & h_{12}^3 \\ h_{11}^4 & h_{12}^4 \\ h_{11}^5 & h_{12}^5 \end{bmatrix} \cdot \begin{bmatrix} \cos 2\theta \\ \sin 2\theta \end{bmatrix}. \quad (1.15)$$

Observe that saying $K_N \neq 0$ is equivalent to say that the rank of the matrix $(h_{1j}^\alpha)_{3 \times 2}$ in (1.15) is two and $\mathbf{B}_p(T_p M)$ is two-dimensional.

The map $\mathbf{g}$ is *superminimal* if in addition, the curvature ellipse is always a circle. That is equivalent to

$$\|\mathbf{B}_{11}\| = \|\mathbf{B}_{12}\|, \quad \langle \mathbf{B}_{11}, \mathbf{B}_{12} \rangle = 0, \quad \mathbf{B}_{ij} := \mathbf{B}(\mathbf{e}_i, \mathbf{e}_j). \quad (1.16)$$

2. Example of a minimal hypersurface with vanishing Gauss-Kronecker curvature

In this section, we shall give an example of a minimal hypersurface in $\mathbb{S}^5$ with vanishing Gauss-Kronecker and 3-mean curvatures. This example is a 2-spherical local bundle over a minimal surface in $\mathbb{S}^5$.

Before to construct the example, we give a result concerning an immersed superminimal surface with vanishing scalar normal curvature.

Proposition 2.1. Let M^2 be a complete surface and $\mathbf{g} : M^2 \hookrightarrow \mathbb{S}^5$ be a superminimal immersion. If the scalar normal curvature K_N vanishes, then M^2 is compact and $\mathbf{g}(M^2)$ is a totally geodesic sphere in $\mathbb{S}^5$.

Proof. It is easy to see that

$$\begin{aligned} (K-1)^2 - \frac{1}{4}K_N &= (\|\mathbf{B}_{11}\|^2 + \|\mathbf{B}_{12}\|^2)^2 - 4 \sum_{\alpha < \beta} \begin{vmatrix} h_{11}^\alpha & h_{12}^\alpha \\ h_{11}^\beta & h_{12}^\beta \end{vmatrix} \\ &= (\|\mathbf{B}_{11}\|^2 + \|\mathbf{B}_{12}\|^2)^2 - 4(\|\mathbf{B}_{11}\|^2 \|\mathbf{B}_{12}\|^2 - \langle \mathbf{B}_{11}, \mathbf{B}_{12} \rangle^2) \\ &= (\|\mathbf{B}_{11}\|^2 - \|\mathbf{B}_{12}\|^2)^2 + 4\langle \mathbf{B}_{11}, \mathbf{B}_{12} \rangle^2. \end{aligned}$$

Therefore, it follows from (1.16) that $\mathbf{g}$ is superminimal if and only if

$$(K-1)^2 - \frac{1}{4}K_N = 0. \quad (2.1)$$

Assume now that $\mathbf{g} : M^2 \hookrightarrow \mathbb{S}^5$ is superminimal and that $K_N = 0$. So, the vectors $\mathbf{B}_{11}$ and $\mathbf{B}_{12}$ are orthogonal and linearly dependent, thus $\mathbf{B} \equiv 0$ and $K = 1$. Therefore, if M^2 is complete, by Bonnet-Myers' Theorem, M^2 is compact and, since $\mathbf{B} \equiv 0$, $\mathbf{g}(M^2)$ is a totally geodesic sphere in $\mathbb{S}^5$. $\square$

Now, starting the construction of the example, let $\mathbf{g}: M^2 \hookrightarrow \mathbb{S}^5$ be an isometric immersion and consider $\mathcal{N} \subset (\mathbf{g}_*(TM))^\perp$ the unit normal bundle of the immersion $\mathbf{g}$. Then

$$\mathcal{N} = \{(p, V) \in M^2 \times \mathbb{R}^6 : \|V\| = 1, V \perp \mathbb{R} \cdot \mathbf{g}(p) \oplus \mathbf{g}_*(T_p M)\}.$$

Denote the projection to the first factor by $\pi_1: \mathcal{N} \rightarrow M^2$ and the projection to the second factor by the map $\mathbf{x}_\mathbf{g}: \mathcal{N} \rightarrow \mathbb{S}^5$.

Consider

$$\mathcal{N}_*(p) = \mathcal{N}(p) \setminus \{\mathbf{B}_p(T_p M)^\perp \cap \mathbf{g}_*(T_p M)^\perp\}. \quad (2.2)$$

We have three situations :

- (a) if $K_N \neq 0$, i.e., if rank of the matrix $(h_{1j}^\alpha)_{3 \times 2}$ is two, then $\mathcal{N}_*(p)$ is a 2-sphere without two antipodal points, given by the orthogonal complement of $\mathbf{B}_p(T_p M)$ in $\mathbf{g}_*(T_p M)^\perp$;
- (b) if rank of the matrix $(h_{1j}^\alpha)_{3 \times 2}$ is one, then $\mathcal{N}_*(p)$ is a 2-sphere without a great circle;
- (c) if rank of the matrix $(h_{1j}^\alpha)_{3 \times 2}$ is zero, that is, if $\mathbf{g}(M)$ is totally geodesic, then $\mathcal{N}_*(p) = \emptyset$.




FIGURE 1. The three cases of $\mathcal{N}_*(p)$ in terms of matrix $(h_{1j}^\alpha)_{3 \times 2}$

Theorem 2.2. *Let $\mathbf{g}: M^2 \hookrightarrow \mathbb{S}^5$ be a minimal immersion with nowhere zero second fundamental form. Then there exists an open set $\mathcal{N}_*$ of $\mathcal{N}$ such that $\mathbf{x}_\mathbf{g}: \mathcal{N}_* \rightarrow \mathbb{S}^5$ is an immersion of the 4-dimensional manifold $\mathcal{N}_*$. Furthermore, $\mathbf{x}_\mathbf{g}: \mathcal{N}_* \rightarrow \mathbb{S}^5$ is an immersed minimal hypersurface of $\mathbb{S}^5$ with zero Gauss-Kronecker curvature and zero 3-mean curvature.*

Proof. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be a frame defined on an open set $U \subset M^2$ adapted to the immersion $\mathbf{g}$ and let W be a coordinate neighborhood of $\mathbb{S}^2$. Parameterize $\pi_1^{-1}(U)$ locally by $U \times W$ via the map $\mathbf{y}: U \times W \rightarrow \mathcal{N}$ given by

$$(p, \theta, \varphi) \mapsto (p, \sin \varphi \cos \theta \mathbf{e}_3 + \sin \varphi \sin \theta \mathbf{e}_4 + \cos \varphi \mathbf{e}_5), \quad 0 \leq \theta < 2\pi, \quad 0 < \varphi < \pi.$$

Then $\mathbf{x}: U \times W \rightarrow \mathbb{S}^5$ given by

$$\mathbf{x}(p, \theta, \varphi) = \sin \varphi \cos \theta \mathbf{e}_3 + \sin \varphi \sin \theta \mathbf{e}_4 + \cos \varphi \mathbf{e}_5 \quad (2.3)$$

is a local representation of $\mathbf{x}_{\mathbf{g}}$. Thus,

$$\begin{aligned} d\mathbf{x} = & \sin \varphi \cos \theta d\mathbf{e}_3 + \sin \varphi \sin \theta d\mathbf{e}_4 + \cos \varphi d\mathbf{e}_5 + \\ & + \cos \varphi \cos \theta d\varphi \mathbf{e}_3 + \cos \varphi \sin \theta d\varphi \mathbf{e}_4 - \sin \varphi d\varphi \mathbf{e}_5 - \\ & - \sin \varphi \sin \theta d\theta \mathbf{e}_3 + \sin \varphi \cos \theta d\theta \mathbf{e}_4 \end{aligned}$$

The structure equations adapted to $\mathbf{g}$ then yield

$$\begin{aligned} d\mathbf{x} = & (\sin \varphi \cos \theta \omega_3^1 + \sin \varphi \sin \theta \omega_4^1 + \cos \varphi \omega_5^1) \otimes \mathbf{e}_1 + \\ & + (\sin \varphi \cos \theta \omega_3^2 + \sin \varphi \sin \theta \omega_4^2 + \cos \varphi \omega_5^2) \otimes \mathbf{e}_2 + \\ & + (-\sin \varphi \sin \theta d\theta + \cos \varphi \cos \theta d\varphi + \cos \varphi \omega_5^3 - \sin \varphi \sin \theta \omega_3^4) \otimes \mathbf{e}_3 + \\ & + (\sin \varphi \cos \theta d\theta + \cos \varphi \sin \theta d\varphi + \cos \varphi \omega_5^4 + \sin \varphi \cos \theta \omega_3^4) \otimes \mathbf{e}_4 + \\ & + (-\sin \varphi d\varphi + \sin \varphi \cos \theta \omega_3^5 + \sin \varphi \sin \theta \omega_4^5) \otimes \mathbf{e}_5. \end{aligned} \quad (2.4)$$

From (1.4) and (2.4), we obtain the first fundamental form of $\mathbf{x}$

$$\begin{aligned} ds_{\mathbf{x}}^2 = \langle d\mathbf{x}, d\mathbf{x} \rangle = & \begin{bmatrix} \omega^1 & \omega^2 \end{bmatrix} \otimes \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} \cdot \begin{bmatrix} \omega^1 \\ \omega^2 \end{bmatrix} + \\ & + (-\sin \varphi \sin \theta d\theta + \cos \varphi \cos \theta d\varphi + \cos \varphi \omega_5^3 - \sin \varphi \sin \theta \omega_3^4)^2 + \\ & + (\sin \varphi \cos \theta d\theta + \cos \varphi \sin \theta d\varphi + \cos \varphi \omega_5^4 + \sin \varphi \cos \theta \omega_3^4)^2 + \\ & + (-\sin \varphi d\varphi + \sin \varphi \cos \theta \omega_3^5 + \sin \varphi \sin \theta \omega_4^5)^2, \end{aligned}$$

where

$$c_{ij} = \sum_k a_{ik} a_{kj}, \quad a_{ij} = \sin \varphi \cos \theta h_{ij}^3 + \sin \varphi \sin \theta h_{ij}^4 + \cos \varphi h_{ij}^5.$$

Since $\mathbf{g}$ is minimal then

$$C = A^2 = \begin{bmatrix} a_{11}^2 + a_{12}^2 & 0 \\ 0 & a_{11}^2 + a_{12}^2 \end{bmatrix} = \begin{bmatrix} -\det A & 0 \\ 0 & -\det A \end{bmatrix}.$$

Thus

$$\begin{aligned} ds_{\mathbf{x}}^2 = & -\det A ((\omega^1)^2 + (\omega^2)^2) + \\ & + (-\sin \varphi \sin \theta d\theta + \cos \varphi \cos \theta d\varphi + \cos \varphi \omega_5^3 - \sin \varphi \sin \theta \omega_3^4)^2 + \\ & + (\sin \varphi \cos \theta d\theta + \cos \varphi \sin \theta d\varphi + \cos \varphi \omega_5^4 + \sin \varphi \cos \theta \omega_3^4)^2 + \\ & + (-\sin \varphi d\varphi + \sin \varphi \cos \theta \omega_3^5 + \sin \varphi \sin \theta \omega_4^5)^2, \end{aligned} \quad (2.5)$$

the quadratic form $-\det A ((\omega^1)^2 + (\omega^2)^2)$ is positive and is positive-definite if and only if $\det C \neq 0$. Since

$$\begin{aligned} \det C = & \left[(\sin \varphi \cos \theta h_{11}^3 + \sin \varphi \sin \theta h_{11}^4 + \cos \varphi h_{11}^5)^2 + \right. \\ & \left. + (\sin \varphi \cos \theta h_{12}^3 + \sin \varphi \sin \theta h_{12}^4 + \cos \varphi h_{12}^5)^2 \right]^2, \end{aligned}$$

we get

$$\det C = 0 \iff \begin{bmatrix} h_{11}^3 & h_{11}^4 & h_{11}^5 \\ h_{12}^3 & h_{12}^4 & h_{12}^5 \end{bmatrix} \cdot \begin{bmatrix} \sin \varphi \cos \theta \\ \sin \varphi \sin \theta \\ \cos \varphi \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \quad (2.6)$$

From the hypothesis that the second fundamental form is nonzero, there are two possibilities at each point p : the rank of $(h_{1j}^\alpha)_{3 \times 2}$ is either two or one. We claim that in both cases $\det C \neq 0$ in all point $(p, \theta, \varphi) \in \mathcal{N}_*(p)$.

- (i) Suppose $\text{rank}((h_{1j}^\alpha)_{3 \times 2}) = 2$. In this case, we may choose a particular orthonormal local frame $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ such that the matrix $(h_{1j}^\alpha)_{3 \times 2}$ takes the form

$$\begin{bmatrix} \alpha & \beta \\ 0 & \gamma \\ 0 & 0 \end{bmatrix},$$

with $\alpha\gamma \neq 0$. In fact, starting from $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$, an orthonormal frame, with generic matrix (h_{1j}^α) , let

$$\begin{aligned} \tilde{\mathbf{e}}_3 &= h_{11}^3 \mathbf{e}_3 + h_{11}^4 \mathbf{e}_4 + h_{11}^5 \mathbf{e}_5 \\ \tilde{\mathbf{e}}_4 &= h_{12}^3 \mathbf{e}_3 + h_{12}^4 \mathbf{e}_4 + h_{12}^5 \mathbf{e}_5. \end{aligned}$$

One may apply the Gram Schmidt process on these two vectors to obtain the vectors $\{\tilde{\mathbf{e}}_3, \tilde{\mathbf{e}}_4\}$. Then we chose $\tilde{\mathbf{e}}_5$, such that $\{\mathbf{e}_1, \mathbf{e}_2, \tilde{\mathbf{e}}_3, \tilde{\mathbf{e}}_4, \tilde{\mathbf{e}}_5\}$ be orthonormal. This is the required basis. It is easy to verify that $\alpha = \|\tilde{\mathbf{e}}_3\|$, $\alpha\beta = \langle \tilde{\mathbf{e}}_3, \tilde{\mathbf{e}}_4 \rangle$ and $\beta^2 + \gamma^2 = \|\tilde{\mathbf{e}}_4\|^2$.

For a such particular frame, the equivalence (2.6) reduces to

$$\det C = 0 \iff \begin{cases} \alpha \sin \varphi \cos \theta &= 0 \\ \beta \sin \varphi \cos \theta + \gamma \sin \varphi \sin \theta &= 0 \end{cases}.$$

Since $\alpha\gamma \neq 0$ and $0 < \varphi < \pi$, then $\det C \neq 0$ for all point $(p, \theta, \varphi) \in \mathcal{N}_*(p)$, given in (2.2).

- (ii) Suppose $\text{rank}((h_{1j}^\alpha)_{3 \times 2}) = 1$. We may choose a particular orthonormal local frame $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ such that the matrix $(h_{1j}^\alpha(p))_{3 \times 2}$, at point p , takes the form

$$\begin{bmatrix} \alpha & \beta \\ 0 & 0 \\ 0 & 0 \end{bmatrix},$$

with $\alpha^2 + \beta^2 \neq 0$. In this case, the system (2.6) takes the form

$$\det C = 0 \iff \begin{cases} \alpha \sin \varphi \cos \theta &= 0 \\ \beta \sin \varphi \cos \theta &= 0 \end{cases}.$$

Since $\alpha^2 + \beta^2 \neq 0$ and $0 < \varphi < \pi$, then $\det C \neq 0$ if and only if θ is different from $\frac{\pi}{2}$ and $\frac{3\pi}{2}$. Then $\det C \neq 0$ for all point $(p, \theta, \varphi) \in \mathcal{N}_*(p)$, given in (2.2).

Let

$$\mathbf{w} = a \mathbf{e}_1 + b \mathbf{e}_2 + c \frac{\partial}{\partial \theta} + d \frac{\partial}{\partial \varphi}, \quad a, b, c, d \in \mathbb{R}$$

be a vector in $T_p U \times T_p W$. The 1-forms ω_β^α , are the connection forms of the normal bundle of $\mathbf{g}$ and $\omega_\beta^\alpha(\frac{\partial}{\partial \theta})$, $\omega_\beta^\alpha(\frac{\partial}{\partial \varphi})$ are zero. Suppose now

$$d\mathbf{x}_{(p, \theta, \varphi)}(\mathbf{w}) = 0.$$

From equation (2.5) and from the facts that $ds_{\mathbf{x}}^2(\mathbf{w}) = 0$ and $\det C \neq 0$, we get $a = b = 0$ and c and d must satisfy

$$\begin{bmatrix} -\sin \varphi \sin \theta & \cos \varphi \cos \theta \\ \sin \varphi \cos \theta & \cos \varphi \sin \theta \\ 0 & -\sin \varphi \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Since $0 < \varphi < \pi$, the first matrix on the equation above has rank 2 and necessarily $c = d = 0$. So,

$$d\mathbf{x}_{(p, \theta, \varphi)}(\mathbf{w}) = 0 \quad \Rightarrow \quad \mathbf{w} = 0.$$

Therefore, $\mathbf{x}_{\mathbf{g}}$ is an immersion.

Let $\{\mathbf{e}_1, \mathbf{e}_2, \frac{\partial}{\partial \theta}, \frac{\partial}{\partial \varphi}, \mathbf{g} \circ \pi_1\}$ an orthogonal frame adapted to the immersion $\mathbf{x}_{\mathbf{g}}$. Note that $\mathbf{g} \circ \pi_1(p, \theta, \varphi)$ is normal to $T_{\mathbf{x}(p, \theta, \varphi)}\mathcal{N}_*$ and tangent to $\mathbb{S}^5$. The second fundamental form of $\mathbf{x}_{\mathbf{g}}$ is given by

$$\begin{aligned} \Pi_{\mathbf{x}_{\mathbf{g}}} &= -\langle d\mathbf{x}, d(\mathbf{g} \circ \pi_1) \rangle \\ &= -\langle d\mathbf{x}, \omega^1 \otimes \mathbf{e}_1 + \omega^2 \otimes \mathbf{e}_2 \rangle \\ &= a_{11} \omega^1 \otimes \omega^1 + 2a_{12} \omega^1 \otimes \omega^2 + a_{22} \omega^2 \otimes \omega^2. \end{aligned}$$

Since $\mathbf{g}$ is minimal, then the trace of matrix $A = (a_{ij})$ is zero. Thus, trace of matrix of the $\Pi_{\mathbf{x}_{\mathbf{g}}}$ is zero too. Therefore $\mathbf{x}_{\mathbf{g}}: \mathcal{N}_* \rightarrow \mathbb{S}^5$ is a minimal immersion. Since the rank of matrix A is two, then 3 and 4-mean curvatures vanish. This proves the theorem. $\square$

3. Minimal hypersurfaces in $\mathbb{S}^5$ with zero Gauss-Kronecker and 3-mean curvatures

In this section we shall study locally the complete minimal hypersurfaces in $\mathbb{S}^5$ with zero Gauss-Kronecker curvature, zero 3-mean curvature and nowhere zero second fundamental form. First we need the following lemmas.

Lemma 3.1. *Let $\mathbf{g}: M^4 \rightarrow \mathbb{S}^5$ be an immersed hypersurface of $\mathbb{S}^5$ and let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be an orthonormal (local) frame adapted to $\mathbf{g}$ with dual co-frame $\{\omega^1, \omega^2, \omega^3, \omega^4, \omega^5\}$. If $\{\mathbf{e}_1, \mathbf{e}_2, \tilde{\mathbf{e}}_3, \tilde{\mathbf{e}}_4, \mathbf{e}_5\}$ is other orthonormal frame with dual co-frame $\{\omega^1, \omega^2, \tilde{\omega}^3, \tilde{\omega}^4, \omega^5\}$ such that*

$$\begin{bmatrix} \tilde{\mathbf{e}}_3 \\ \tilde{\mathbf{e}}_4 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \mathbf{e}_3 \\ \mathbf{e}_4 \end{bmatrix}.$$

Then

$$\begin{aligned} \begin{bmatrix} \tilde{\omega}^3 \\ \tilde{\omega}^4 \end{bmatrix} &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \omega^3 \\ \omega^4 \end{bmatrix}; \\ \begin{bmatrix} \tilde{\omega}_3^1 & \tilde{\omega}_3^2 \\ \tilde{\omega}_4^1 & \tilde{\omega}_4^2 \end{bmatrix} &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \omega_3^1 & \omega_3^2 \\ \omega_4^1 & \omega_4^2 \end{bmatrix}; \\ \tilde{\omega}_2^1 &= \omega_2^1, \quad \tilde{\omega}_4^3 = \omega_4^3 + d\theta, \quad \tilde{\omega}_i^5 = \omega_i^5. \end{aligned}$$

Proof. Taking exterior differentiate of

$$\begin{cases} \omega^3 = \cos \theta \tilde{\omega}^3 + \sin \theta \tilde{\omega}^4, \\ \omega^4 = -\sin \theta \tilde{\omega}^3 + \cos \theta \tilde{\omega}^4 \end{cases}$$

and using (1.8) we obtain

$$\begin{aligned} d\tilde{\omega}^3 &= -(\cos \theta \omega_1^3 - \sin \theta \omega_1^4) \wedge \omega^1 - (\cos \theta \omega_2^3 - \sin \theta \omega_2^4) \wedge \omega^2 - (\omega_4^3 + d\theta) \wedge \tilde{\omega}^4 \\ d\tilde{\omega}^4 &= -(\sin \theta \omega_1^3 + \cos \theta \omega_1^4) \wedge \omega^1 - (\sin \theta \omega_2^3 + \cos \theta \omega_2^4) \wedge \omega^2 + (\omega_4^3 + d\theta) \wedge \tilde{\omega}^3. \end{aligned}$$

□

Lemma 3.2. *In the hypothesis above. If the functions f_3, f_4, g_3 and g_4 are defined by*

$$\begin{cases} \omega_2^3 = f_3 \omega^1 + g_3 \omega^2 \\ \omega_2^4 = f_4 \omega^1 + g_4 \omega^2. \end{cases} \quad (3.1)$$

Then

$$\begin{bmatrix} \tilde{f}_3 & \tilde{g}_3 \\ \tilde{f}_4 & \tilde{g}_4 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} f_3 & g_3 \\ f_4 & g_4 \end{bmatrix}$$

and

$$\begin{aligned} \tilde{f}_3^2 + \tilde{f}_4^2 &= f_3^2 + f_4^2, & \tilde{f}_3 \tilde{g}_3 + \tilde{f}_4 \tilde{g}_4 &= f_3 g_3 + f_4 g_4, \\ \tilde{g}_3^2 + \tilde{g}_4^2 &= g_3^2 + g_4^2, & \tilde{f}_3 \tilde{g}_4 - \tilde{f}_4 \tilde{g}_3 &= f_3 g_4 - f_4 g_3. \end{aligned}$$

Proof. Since

$$\begin{cases} \mathbf{e}_3 = \cos \theta \tilde{\mathbf{e}}_3 + \sin \theta \tilde{\mathbf{e}}_4, \\ \mathbf{e}_4 = -\sin \theta \tilde{\mathbf{e}}_3 + \cos \theta \tilde{\mathbf{e}}_4. \end{cases}$$

It follows from (3.1) that

$$\begin{aligned} (\nabla \mathbf{e}_2) &= \omega_2^1 \otimes \mathbf{e}_1 + \omega_2^3 \otimes \mathbf{e}_3 + \omega_2^4 \otimes \mathbf{e}_4 = \\ &= \omega_2^1 \otimes \mathbf{e}_1 + [(\cos \theta f_3 - \sin \theta f_4) \omega^1 + (\cos \theta g_3 - \sin \theta g_4) \omega^2] \otimes \tilde{\mathbf{e}}_3 + \\ &\quad + [(\sin \theta f_3 + \cos \theta f_4) \omega^1 + (\sin \theta g_3 + \cos \theta g_4) \omega^2] \otimes \tilde{\mathbf{e}}_4, \end{aligned}$$

where ∇ is the Levi-Civita connections of M^4 .

□

Theorem 3.3. *Let $\mathbf{g} : M^4 \rightarrow \mathbb{S}^5$ be a complete, oriented minimal immersed hypersurface of $\mathbb{S}^5$ with zero Gauss-Kronecker curvature and zero 3-mean curvature. If the square S of the length of the second fundamental form is nowhere zero, then there exists a minimal immersed $\tilde{\eta} : \mathbb{V}^2 \rightarrow \mathbb{S}^5$ and a local isometry $\tau : M_*^4 \rightarrow \mathcal{N}_*$, where $M_*^4 = M^4 \setminus \{\mathbf{g}^{-1}(\mathbf{B}_{\pi(p)}(T_{\pi(p)}\mathbb{V})^\perp \cap \tilde{\eta}_*(T_{\pi(p)}\mathbb{V})^\perp)\}$, such that $\eta|_{M_*^4} = \tilde{\eta} \circ \pi_1 \circ \tau$ and $\mathbf{x}_{\tilde{\eta}} \circ \tau = \mathbf{g}|_{M_*^4}$.*

Proof. Let $\mathcal{B} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be an orthonormal frame on $\mathbb{S}^5$ adapted to the immersion $\mathbf{g} : M^4 \rightarrow \mathbb{S}^5$ in the sense that $\{\mathbf{e}_1, \dots, \mathbf{e}_4\}$ spans $\mathbf{g}_*(TM)$, $\mathbf{e}_5$ determines a global Gauss map $\eta : M^4 \rightarrow \mathbb{S}^5$ and as described in (1.7) holds

$$\omega_i^5 = \lambda_i \omega^i, \quad i = 1, 2, 3, 4.$$

By hypothesis and (1.10), we have exactly two of the λ_i are zero and the sum of the other two is zero. Hence, we may assume that

$$\lambda = \lambda_1 = -\lambda_2 > 0, \quad \lambda_3 = \lambda_4 = 0$$

and

$$\omega_1^5 = \lambda \omega^1, \quad \omega_2^5 = -\lambda \omega^2, \quad \omega_3^5 = \omega_4^5 = 0. \quad (3.2)$$

In this case, we have two directions and one plane well defined. Since $S > 0$ on M^4 then

$$\mathcal{D}_p := \{v \in T_p M^4 : \mathbf{B}_p(v, w) = 0, \quad \forall w \in T_p M^4\}$$

is a 2-dimensional distribution. Now, we proceed to show that $\mathcal{D}_p$ is involutive, that its integral surfaces $\mathcal{F}^2$ are totally geodesic in M^4 and $\mathbf{g}(\mathcal{F}^2)$ are totally geodesic in $\mathbb{S}^5$. From (3.2) and Codazzi equations given in (1.9) we have that

$$\begin{cases} \omega_1^3 \wedge \omega^1 - \omega_2^3 \wedge \omega^2 = 0; \\ \omega_1^4 \wedge \omega^1 - \omega_2^4 \wedge \omega^2 = 0. \end{cases} \quad (3.3)$$

Let $\mathfrak{I}(\omega^1, \omega^2)$ be the ideal generated by ω^1 and ω^2 . By Cartan's lemma $\omega_1^3, \omega_2^3, \omega_1^4$ and ω_2^4 belong to the ideal $\mathfrak{I}(\omega^1, \omega^2)$. From (1.8) follows

$$d\omega^i \in \mathfrak{I}(\omega^1, \omega^2), \quad i = 1, 2 \quad \text{or} \quad \omega^i([\mathbf{e}_3, \mathbf{e}_4]) = 0, \quad i = 1, 2.$$

Therefore, $\mathcal{D}_p$ is involutive. Then by Frobenius Theorem there exists an unique maximal, connected, integral surface $\mathcal{F}^2$ of $\mathcal{D}_p$ through p . Since $\mathbf{g} : M^4 \rightarrow \mathbb{S}^5$ is an immersion of a complete manifold, a theorem of D. Ferus [4] implies that $\mathcal{F}^2$ is complete and totally geodesic in M^4 . From (3.3) it is easy to see that

$$\begin{cases} \omega_2^3(\mathbf{e}_1) + \omega_1^3(\mathbf{e}_2) = 0; \\ \omega_2^4(\mathbf{e}_1) + \omega_1^4(\mathbf{e}_2) = 0; \\ \omega_j^i(\mathbf{e}_k) = 0, \quad i = 1, 2; \quad j, k = 3, 4. \end{cases} \quad (3.4)$$

Substituting (3.2) and (3.4) into (1.8) we obtain

$$\begin{cases} d\mathbf{e}_3(\mathbf{e}_3) = -\mathbf{g} + \omega_3^4(\mathbf{e}_3)\mathbf{e}_4; \\ d\mathbf{e}_3(\mathbf{e}_4) = \omega_3^4(\mathbf{e}_4)\mathbf{e}_4; \\ d\mathbf{e}_4(\mathbf{e}_3) = \omega_4^3(\mathbf{e}_3)\mathbf{e}_3; \\ d\mathbf{e}_4(\mathbf{e}_4) = -\mathbf{g} + \omega_4^3(\mathbf{e}_4)\mathbf{e}_3. \end{cases} \quad (3.5)$$

This implies that

$$\omega^k(\bar{\nabla}_{\mathbf{e}_j}\mathbf{e}_i) = 0, \quad k = 1, 2; \quad i, j = 3, 4,$$

where $\bar{\nabla}$ is the Levi-Civita connections of $\mathbb{S}^5$. Therefore, $\mathbf{g}(\mathcal{F}^2)$ is totally geodesic in $\mathbb{S}^5$. Since $\mathcal{F}^2$ is complete and $\mathbf{g}|_{\mathcal{F}^2} : \mathcal{F}^2 \rightarrow \mathbb{S}^5$ is an isometric immersion, then $\mathbf{g}(\mathcal{F}^2)$ is an unitary 2-sphere $\mathbb{S}^2$ in $\mathbb{S}^5$. So, we have that $\mathbf{g}|_{\mathcal{F}^2} : \mathcal{F}^2 \rightarrow \mathbb{S}^2$ is a covering map (see [6, p.146, Prop.6.16]). Thus, $\mathcal{F}^2$ is a 2-sphere in M^4 and $\mathbf{g}|_{\mathcal{F}^2} : \mathcal{F}^2 \rightarrow \mathbb{S}^2$ is a diffeomorphism (see [6, p.141]). Therefore, the maximal integral surface $\mathcal{F}^2$ is regular (see [9, p.98]) and a theorem of R. Palais (see [7]) implies that the quotient space

$$\mathbb{V}^2 = M^4 / \mathcal{F}^2$$

can be endowed with a structure of a 2-dimensional differential manifold such that $\pi : M^4 \rightarrow \mathbb{V}^2$ is a submersion. The Gauss map $\eta : M^4 \rightarrow \mathbb{S}^5$ induces a smooth map $\tilde{\eta} : \mathbb{V}^2 \rightarrow \mathbb{S}^5$ such that $\tilde{\eta} \circ \pi = \eta$,

$$\begin{array}{ccc} M^4 & & \\ \pi \downarrow & \searrow \eta & \\ \mathbb{V}^2 & \xrightarrow{\tilde{\eta}} & \mathbb{S}^5. \end{array}$$

In fact, from (3.2), we have that $\bar{\nabla}_{\mathbf{e}_3}\eta$ and $\bar{\nabla}_{\mathbf{e}_4}\eta$ vanish. Thus, η is constant along the integral surfaces $\mathcal{F}^2$ and $\tilde{\eta}$ is well defined. Let S be a smooth transversal surface to the leaf $\mathcal{F}^2$ of $\mathcal{D}_p$ through a point $p \in M^4$ such that $T_p S = \text{span}\{\mathbf{e}_1|_p, \mathbf{e}_2|_p\}$ and $T_p \mathcal{F} = \text{span}\{\mathbf{e}_3|_p, \mathbf{e}_4|_p\}$. Since $\pi : M^4 \rightarrow \mathbb{V}^2$ is a submersion, then

$$\text{span}\{d\pi_p(\mathbf{e}_1|_p), d\pi_p(\mathbf{e}_2|_p)\} = T_{\pi(p)}\mathbb{V}^2.$$

The third formula given in (1.8) implies that

$$\begin{aligned} d\tilde{\eta}_{\pi(p)}(d\pi_p(\mathbf{e}_1|_p)) &= d(\tilde{\eta} \circ \pi)_p(\mathbf{e}_1|_p) = d\eta_p(\mathbf{e}_1|_p) = -\lambda(p)\mathbf{e}_1|_p; \\ d\tilde{\eta}_{\pi(p)}(d\pi_p(\mathbf{e}_2|_p)) &= \lambda(p)\mathbf{e}_2|_p. \end{aligned}$$

So, the first fundamental form of $\tilde{\eta}$ is given by

$$ds_{\tilde{\eta}}^2 = \lambda^2 (\omega^1 \otimes \omega^1 + \omega^2 \otimes \omega^2).$$

Since $\lambda(p) > 0$ for all $p \in M^4$, then $\tilde{\eta} : \mathbb{V}^2 \rightarrow \mathbb{S}^5$ is an immersion. Now, we will show that $\tilde{\eta}$ is a minimal immersion. Let

$$\{X_1, X_2\} = \left\{ d\pi_p \left(\frac{1}{\lambda} \mathbf{e}_1|_p \right), d\pi_p \left(\frac{1}{\lambda} \mathbf{e}_2|_p \right) \right\}$$

be an orthogonal basis for $T_{\pi(p)}\mathbb{V}^2$ and let $\{X_3, X_4, X_5\}$ be an orthogonal frame in the normal bundle of $\tilde{\eta}$ such that

$$X_3 \circ \pi|_S = \mathbf{e}_3|_S, \quad X_4 \circ \pi|_S = \mathbf{e}_4|_S, \quad X_5 \circ \pi|_S = \mathbf{g}|_S.$$

The second fundamental form $\mathbf{B}_{\pi(p)}$ of $\tilde{\eta}$ is determined by the bilinear forms

$$\Pi_{\tilde{\eta}}^{\alpha} := -\langle d\tilde{\eta}, dX_{\alpha} \rangle_{\pi(p)}, \quad \alpha = 3, 4, 5.$$

From (1.8) it is easy to see that

$$\begin{aligned} \Pi_{\tilde{\eta}}^3(X_i \otimes X_j) &= -\frac{1}{\lambda^2} \langle d\eta, d\mathbf{e}_3 \rangle (\mathbf{e}_i|_p \otimes \mathbf{e}_j|_p), \quad i, j = 1, 2; \\ &= \frac{1}{\lambda} (\omega^1 \otimes \omega_3^1 - \omega^2 \otimes \omega_3^2) (\mathbf{e}_i|_p \otimes \mathbf{e}_j|_p); \\ \Pi_{\tilde{\eta}}^4(X_i \otimes X_j) &= \frac{1}{\lambda} (\omega^1 \otimes \omega_4^1 - \omega^2 \otimes \omega_4^2) (\mathbf{e}_i|_p \otimes \mathbf{e}_j|_p); \\ \Pi_{\tilde{\eta}}^5(X_i \otimes X_j) &= \frac{1}{\lambda} (\omega^1 \otimes \omega^1 - \omega^2 \otimes \omega^2) (\mathbf{e}_i|_p \otimes \mathbf{e}_j|_p). \end{aligned}$$

Denote by $\tilde{A}^{\alpha}$ the shape operators of $\tilde{\eta}$ at $\pi(p)$, then

$$\begin{aligned} \tilde{A}^3 &= \frac{1}{\lambda(p)} \begin{bmatrix} \omega_3^1(\mathbf{e}_1) & \omega_3^1(\mathbf{e}_2) \\ -\omega_3^2(\mathbf{e}_1) & -\omega_3^2(\mathbf{e}_2) \end{bmatrix}, \quad \tilde{A}^4 = \frac{1}{\lambda(p)} \begin{bmatrix} \omega_4^1(\mathbf{e}_1) & \omega_4^1(\mathbf{e}_2) \\ -\omega_4^2(\mathbf{e}_1) & -\omega_4^2(\mathbf{e}_2) \end{bmatrix}, \\ \tilde{A}^5 &= \frac{1}{\lambda(p)} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \end{aligned}$$

Hence, $\tilde{\eta}$ is a minimal immersion if and only if

$$\begin{cases} \omega_3^1(\mathbf{e}_1) - \omega_3^2(\mathbf{e}_2) = 0 \\ \omega_4^1(\mathbf{e}_1) - \omega_4^2(\mathbf{e}_2) = 0. \end{cases} \quad (3.6)$$

By taking exterior differentiations of (3.2) we have

$$d\omega_i^5 = d\lambda_i \wedge \omega^i - \lambda_i \omega_k^i \wedge \omega^k.$$

From Codazzi equation given in (1.9) it follows that

$$d\lambda_i \wedge \omega^i + (\lambda_k - \lambda_i) \omega_k^i \wedge \omega^k = 0, \quad i = 1, 2. \quad (3.7)$$

Evaluating these equation on $\mathbf{e}_k \otimes \mathbf{e}_i$ we obtain

$$\begin{cases} \mathbf{e}_1[\lambda] - 2\lambda\omega_2^1(\mathbf{e}_2) = 0 \\ \mathbf{e}_2[\lambda] + 2\lambda\omega_2^1(\mathbf{e}_1) = 0 \end{cases}, \quad (3.8)$$

$$\begin{cases} \mathbf{e}_3[\lambda] + \lambda\omega_3^1(\mathbf{e}_1) = 0 \\ \mathbf{e}_3[\lambda] + \lambda\omega_3^2(\mathbf{e}_2) = 0 \end{cases}, \quad \begin{cases} \mathbf{e}_4[\lambda] + \lambda\omega_4^1(\mathbf{e}_1) = 0 \\ \mathbf{e}_4[\lambda] + \lambda\omega_4^2(\mathbf{e}_2) = 0 \end{cases}, \quad (3.9)$$

$$\begin{cases} 2\omega_2^1(\mathbf{e}_3) + \omega_3^2(\mathbf{e}_1) = 0 \\ 2\omega_2^1(\mathbf{e}_4) + \omega_4^2(\mathbf{e}_1) = 0 \end{cases}. \quad (3.10)$$

Therefore, the equations (3.9) imply that the condition (3.6) is satisfied. Thus,

$$\tilde{A}^3 = \frac{1}{\lambda(p)} \begin{bmatrix} -g_3 & f_3 \\ f_3 & g_3 \end{bmatrix}, \quad \tilde{A}^4 = \frac{1}{\lambda(p)} \begin{bmatrix} -g_4 & f_4 \\ f_4 & g_4 \end{bmatrix},$$

$$\tilde{A}^5 = \frac{1}{\lambda(p)} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

From (3.4), (3.8), (3.9) and (3.10), we have the skew-symmetric matrices

$$\begin{aligned} (\omega_j^i(\mathbf{e}_1)) &= \begin{bmatrix} 0 & -\frac{1}{2}\mathbf{e}_2[\ln \lambda] & -g_3 & -g_4 \\ * & 0 & -f_3 & -f_4 \\ * & * & 0 & \omega_4^3(\mathbf{e}_1) \\ * & * & * & 0 \end{bmatrix}, \\ (\omega_j^i(\mathbf{e}_2)) &= \begin{bmatrix} 0 & \frac{1}{2}\mathbf{e}_1[\ln \lambda] & f_3 & f_4 \\ * & 0 & -g_3 & -g_4 \\ * & * & 0 & \omega_4^3(\mathbf{e}_2) \\ * & * & * & 0 \end{bmatrix}, \\ (\omega_j^i(\mathbf{e}_3)) &= \begin{bmatrix} 0 & \frac{f_3}{2} & 0 & 0 \\ * & 0 & 0 & 0 \\ * & * & 0 & \omega_4^3(\mathbf{e}_3) \\ * & * & * & 0 \end{bmatrix}, \quad (\omega_j^i(\mathbf{e}_4)) = \begin{bmatrix} 0 & \frac{f_4}{2} & 0 & 0 \\ * & 0 & 0 & 0 \\ * & * & 0 & \omega_4^3(\mathbf{e}_4) \\ * & * & * & 0 \end{bmatrix}, \end{aligned}$$

and relations

$$\begin{cases} \mathbf{e}_3[\lambda] = \lambda g_3 \\ \mathbf{e}_4[\lambda] = \lambda g_4. \end{cases} \quad (3.11)$$

Using the equations given in (1.13) we have that K and K_N of the minimal immersion $\tilde{\eta}: \mathbb{V}^2 \rightarrow \mathbb{S}^5$ are given by

$$K(\pi(p)) = 1 - \frac{1}{\lambda^2} (1 + f_3^2 + f_4^2 + g_3^2 + g_4^2), \quad (3.12)$$

$$R_{412}^3 = \frac{2}{\lambda^2} (f_3 g_4 - f_4 g_3), \quad R_{512}^3 = -\frac{2f_3}{\lambda^2}, \quad R_{512}^4 = -\frac{2f_4}{\lambda^2}, \quad (3.13)$$

$$K_N(\pi(p)) = \frac{16}{\lambda^4} (f_3^2 + f_4^2 + (f_3 g_4 - f_4 g_3)^2). \quad (3.14)$$

By a straightforward calculation it follows that

$$\begin{aligned} [\mathbf{e}_1, \mathbf{e}_2] &= -\frac{1}{2}\mathbf{e}_2[\ln \lambda]\mathbf{e}_1 + \frac{1}{2}\mathbf{e}_1[\ln \lambda]\mathbf{e}_2 + 2f_3\mathbf{e}_3 + 2f_4\mathbf{e}_4, \\ [\mathbf{e}_3, \mathbf{e}_1] &= g_3\mathbf{e}_1 + \frac{f_3}{2}\mathbf{e}_2 + \omega_4^3(\mathbf{e}_1)\mathbf{e}_4, \\ [\mathbf{e}_3, \mathbf{e}_2] &= -\frac{f_3}{2}\mathbf{e}_1 + g_3\mathbf{e}_2 + \omega_4^3(\mathbf{e}_2)\mathbf{e}_4, \\ [\mathbf{e}_4, \mathbf{e}_1] &= g_4\mathbf{e}_1 + \frac{f_4}{2}\mathbf{e}_2 - \omega_4^3(\mathbf{e}_1)\mathbf{e}_3, \\ [\mathbf{e}_4, \mathbf{e}_2] &= -\frac{f_4}{2}\mathbf{e}_1 + g_4\mathbf{e}_2 - \omega_4^3(\mathbf{e}_2)\mathbf{e}_3, \\ [\mathbf{e}_3, \mathbf{e}_4] &= \omega_4^3(\mathbf{e}_3)\mathbf{e}_3 + \omega_4^3(\mathbf{e}_4)\mathbf{e}_4. \end{aligned} \quad (3.15)$$

Applying the relation between the bracket operation on vector fields and the exterior differentiation of 1-form, we obtain the derivatives of the functions f_i and g_i :

$$\begin{aligned} \mathbf{e}_1[f_3] &= -\mathbf{e}_2[g_3] - \omega_4^3(\mathbf{e}_1)f_4 - \omega_4^3(\mathbf{e}_2)g_4 \\ \mathbf{e}_2[f_3] &= \mathbf{e}_1[g_3] + \omega_4^3(\mathbf{e}_1)g_4 - \omega_4^3(\mathbf{e}_2)f_4 \\ \mathbf{e}_1[f_4] &= -\mathbf{e}_2[g_4] + \omega_4^3(\mathbf{e}_1)f_3 + \omega_4^3(\mathbf{e}_2)g_3 \\ \mathbf{e}_2[f_4] &= \mathbf{e}_1[g_4] - \omega_4^3(\mathbf{e}_1)g_3 + \omega_4^3(\mathbf{e}_2)f_3; \end{aligned} \quad (3.16)$$

$$\begin{aligned} \mathbf{e}_3[f_3] &= 2f_3g_3 - \omega_4^3(\mathbf{e}_3)f_4 \\ \mathbf{e}_3[f_4] &= f_3g_4 + f_4g_3 + \omega_4^3(\mathbf{e}_3)f_3 \\ \mathbf{e}_3[g_3] &= g_3^2 - f_3^2 + 1 - \omega_4^3(\mathbf{e}_3)g_4 \\ \mathbf{e}_3[g_4] &= g_3g_4 - f_3f_4 + \omega_4^3(\mathbf{e}_3)g_3; \end{aligned} \quad (3.17)$$

$$\begin{aligned} \mathbf{e}_4[f_3] &= f_3g_4 + f_4g_3 - \omega_4^3(\mathbf{e}_4)f_4 \\ \mathbf{e}_4[f_4] &= 2f_4g_4 + \omega_4^3(\mathbf{e}_4)f_3 \\ \mathbf{e}_4[g_3] &= g_3g_4 - f_3f_4 - \omega_4^3(\mathbf{e}_4)g_4 \\ \mathbf{e}_4[g_4] &= g_4^2 - f_4^2 + 1 + \omega_4^3(\mathbf{e}_4)g_3. \end{aligned} \quad (3.18)$$

By a straightforward calculation, using (3.17) and (3.18), we have that

$$\begin{aligned} \mathbf{e}_3[f_3^2 + f_4^2 + g_3^2 + g_4^2] &= 2g_3(f_3^2 + f_4^2 + g_3^2 + g_4^2 + 1) \\ \mathbf{e}_4[f_3^2 + f_4^2 + g_3^2 + g_4^2] &= 2g_4(f_3^2 + f_4^2 + g_3^2 + g_4^2 + 1) \\ \mathbf{e}_3[R_{412}^3] &= R_{512}^4 \\ \mathbf{e}_4[R_{412}^3] &= -R_{512}^3 \\ \mathbf{e}_3[R_{512}^3] &= -\omega_4^3(\mathbf{e}_3)R_{512}^4 \\ \mathbf{e}_4[R_{512}^3] &= R_{412}^3 - \omega_4^3(\mathbf{e}_4)R_{512}^4 \\ \mathbf{e}_3[R_{512}^4] &= -R_{412}^3 + \omega_4^3(\mathbf{e}_3)R_{512}^3 \\ \mathbf{e}_4[R_{512}^4] &= \omega_4^3(\mathbf{e}_4)R_{512}^3. \end{aligned} \quad (3.19)$$

Using the above equations it follows that

$$\mathbf{e}_3[\mathbf{K}] = 0, \quad \mathbf{e}_4[\mathbf{K}] = 0, \quad \mathbf{e}_3[\mathbf{K}_N] = 0, \quad \mathbf{e}_4[\mathbf{K}_N] = 0.$$

Therefore, the functions $\mathbf{K}$ and $\mathbf{K}_N$ are constant on $\mathcal{F}^2$, which is according with the fact that $\tilde{\eta}$ is well defined.

By Theorem 2.2 $\mathbf{x}_{\tilde{\eta}} : \mathcal{N}_* \rightarrow \mathbb{S}^5$ is an immersed minimal hypersurface of $\mathbb{S}^5$ with zero Gauss-Kronecker and 3-mean curvatures.

Define the map $\tau : M_*^4 \rightarrow \mathcal{N}_*$ by $\tau(p) = (\pi(p), \mathbf{g}(p))$. Thus $\mathbf{x}_{\tilde{\eta}} \circ \tau(p) = \mathbf{g}(p)$. Since, $\pi = \pi_1 \circ \tau$ we have that $\eta|_{M_*^4} = \tilde{\eta} \circ \pi_1 \circ \tau$. Finally, the metric on $\mathcal{N}_*$ is induced by $\mathbf{x}_{\tilde{\eta}}$, then τ must be a local isometry. This proves the theorem. $\square$

References

1. A. C. Asperti, R. M. B. Chaves, and L. A. M. Sousa, Jr., *The Gauss-Kronecker curvature of minimal hypersurfaces in four-dimensional space forms*, Math. Z. **267** (2011), no. 3-4, 523–533. MR 2776046
2. Marcos Dajczer and Detlef Gromoll, *Gauss parametrizations and rigidity aspects of submanifolds*, J. Differential Geom. **22** (1985), no. 1, 1–12. MR **826420** (87g:53088a)
3. S. C. de Almeida and F. G. B. Brito, *Minimal hypersurfaces of S^4 with constant Gauss-Kronecker curvature*, Math. Z. **195** (1987), no. 1, 99–107. MR **888131** (88i:53095)
4. D. Ferus, *On the completeness of nullity foliations.*, The Michigan Mathematical Journal **18** (1971), no. 1, 61–64.
5. Th. Hasanis, A. Savas-Halilaj, and Th. Vlachos, *Complete minimal hypersurfaces in S^4 with zero Gauss-Kronecker curvature*, Math. Proc. Cambridge Philos. Soc. **142** (2007), no. 1, 125–132. MR **2296396** (2007m:53077)
6. Elon Lages Lima, *Fundamental groups and covering spaces*, A K Peters Ltd., Natick, MA, 2003, Translated from the Portuguese by Jonas Gomes. MR **2000701** (2005c:57001)
7. Richard S. Palais, *A global formulation of the Lie theory of transformation groups*, Mem. Amer. Math. Soc. **No. 22** (1957), no. 22, iii–123. MR 0121424 (22 #12162)
8. Jayakumar Ramanathan, *Minimal hypersurfaces in S^4 with vanishing Gauss-Kronecker curvature*, Math. Z. **205** (1990), no. 4, 645–658. MR **1082881** (91m:53048)
9. Itiro Tamura, *Topology of foliations: an introduction*, Translations of Mathematical Monographs, vol. 97, American Mathematical Society, Providence, RI, 1992, Translated from the 1976 Japanese edition and with an afterword by Kiki Hudson, With a foreword by Takashi Tsuboi. MR **1151624** (93c:57021)

Marcos M. Diniz
 Universidade Federal do Pará
 Instituto de Ciências Exatas e Naturais
 e-mail: mdiniz@ufpa.br

José Antonio M. Vilhena
 Universidade Federal do Pará
 Instituto de Ciências Exatas e Naturais
 e-mail: vilhena@ufpa.br

Juan F. Z. Zapata
 Universidade de São Paulo
 Instituto de Matemática e Estatística
 e-mail: jfzzmat@ime.usp.br